

**RESEARCH****Open Access**

Validation of *Aedes aegypti* Aag-2 cells as a model for insect immune studies

Ana Beatriz Ferreira Barletta^{1,2}, Maria Clara L Nascimento Silva³ and Marcos H Ferreira Sorgine^{1,2*}**Abstract**

Background: The understanding of mosquito immune responses can provide valuable tools for development of novel mosquito control strategies. Aiming the study at insect innate immunity, continuous insect cell lines have been established and used as research tools due to the fact that they constitute more homogeneous, sensitive, and reproducible systems than the insects from which they originated. More recently, Aag-2, an *Aedes aegypti* cell lineage, began to be frequently used as a model for studies of mosquito immunity. Nevertheless, to our knowledge, no study has systematically characterized the responses of Aag-2 cell line against different kinds of pathogens and compared its response to those exhibited by whole mosquitoes. For this reason, in this study we characterized gene expression profiles of the Aag-2 cell line in response to different kinds of immune challenges, such as Gram negative and positive bacteria, fungi and viruses, comparing the obtained results with the ones already described in the literature for whole mosquitoes.

Methods: *Aedes aegypti* Aag-2 cells were exposed to different immune stimuli (gram-positive and gram negative heat inactivated bacteria, zymosan or Sindbis virus) for 24 hours and the expression of selected marker genes from toll, IMD and Jak/STAT pathways was analyzed by qPCR. Also, cells were incubated with fluorescent latex beads for evaluation of its phagocytosis capacity.

Results: Aag-2 cells were stimulated with two concentrations of heat-killed Gram negative (*Enterobacter cloacae*) or Gram positive (*Micrococcus luteus*) bacteria, Zymosan or infected with Sindbis virus and the expression of key genes from the main immune related pathways, Toll, IMD and Jak/STAT, were investigated. Our results suggest that Toll and IMD pathways are activated in response to both Gram positive and negative bacteria and Zymosan in Aag-2 cells, displaying an immune profile similar to those described in the literature for whole mosquitoes. The same stimuli were also capable of activating Jak/STAT pathway in Aag-2 cells. Infection with Sindbis virus led to an up-regulation of the transcription factor STAT but was not able to induce the expression of any other gene from any of the pathways assayed. We also showed that this cell line is able to phagocytose latex beads in culture.

Conclusions: Our results characterize the expression profile of Aag-2 cells in response to different immune stimuli and demonstrate that this cell lineage is immune-competent and closely resembles the response described for whole *Ae. aegypti* mosquitoes. Hence, our findings support the use of Aag-2 as a tool to comprehend *Ae. aegypti* immune response both at cellular and humoral levels.

* Correspondence: sorgine@bioqmed.ufrj.br¹Laboratório de Bioquímica de Artrópodes Hematófagos, Instituto de Bioquímica Médica, Programa de Biologia Molecular e Biotecnologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil²Instituto Nacional de Ciência e Tecnologia em Entomologia Molecular (INCT-EM), Rio de Janeiro, Brazil

Full list of author information is available at the end of the article

Background

Aedes aegypti mosquitoes are important vectors of viral diseases, such as Yellow and Dengue fever, which have significant impact on human morbidity and mortality. The incidence of Dengue has grown around the world, mainly in the tropics and subtropics, in the last decades. Nowadays, about 2.5 billion people are currently at risk of infection [1]. Traditional strategies currently employed for the control of the disease vector, *Ae. aegypti*, such as insecticide applications are becoming more and more ineffective due to the rapid development of resistance by this vector [2]. This situation reinforces the need to understand the biology of the vector/pathogen interaction. One aspect that plays an important role in the interaction between a pathogen and its host is immunity. Insect innate immune responses are controlled by three major signaling pathways, the Toll, the Immune Deficiency (IMD) and the Janus kinase (Jak) – signal transducer and activator of transcription (STAT) pathways [3-7].

The Toll pathway is activated mainly by Gram-positive bacteria, fungi [8,9] and viruses [10], and largely controls the expression of antimicrobial peptides (AMPs). When pathogen-associated molecular patterns (PAMPs) are recognized, a signaling cascade is activated and NF- κ B factor(s) is/are translocated to the nucleus to initiate the transcription of molecules involved with the immune response [11]. In *Drosophila melanogaster*, DIF and Dorsal are NF- κ B homologues factors involved in the activation of the Toll pathway, although in mosquitoes only one Toll related NF- κ B factor is present, REL 1, which is homologous to dorsal.

Upon activation of IMD pathway, mainly by Gram negative bacteria in *D. melanogaster*, another NF- κ B factor, Relish, is translocated to the nucleus leading to the transcription of the pathway effector molecules [12,13].

Since the sequencing of the *Ae. aegypti* genome in 2007 [14] putative orthologs of *D. melanogaster* and *An. gambiae* immune genes have been identified and there has been an important growth of the knowledge about how this mosquito is able to fight against different pathogens such as viruses, bacteria and worms [15].

Continuous insect cell lines have been an important research tool for insect biologists since Grace developed the first reported insect cell line from a moth [16]. Cell lineages constitute homogeneous, sensitive and reproducible systems, allowing the detection of very subtle changes in the response to different kinds of pathogens and stimuli. For these reasons, cells from *D. melanogaster* (especially S2 cells), *An. gambiae* (5.1* and Sua5B), *Lutzomyia longipalpis* (LL5 cells), *Ae. albopictus* (especially C6/36 cells), among other insects, have long been used to investigate different aspects of insect immunity, being essential tools for the construction of the

knowledge we have today regarding Toll, IMD, Jak/STAT and RNAi pathways in insects [17-24].

Most of the studies that have focused on mosquito immune responses against pathogens have employed the lineage C6/36 [17,25-27], a lineage established from *Ae. albopictus* larvae homogenates by Singh [28]. C6/36 has been used to understand the regulation of the synthesis and secretion of several important proteins/processes, such as defensin [29], cecropin [24] or phagocytosis [19] but mostly this cell has been used to study aspects of insect-virus relations [17]. This is because one of the main features of C6/36 is the capacity of growing a very broad spectrum of viruses and producing higher virus titers than any other cell [30]. It is now known that this property is, at least partially, due to the lack of a functional RNAi response in these cell lineages [30].

Recently, Aag-2, an *Ae. aegypti* cell lineage of embryonic origin [31] began to be used as a model for studies of mosquito immunity [32-35]. Nevertheless, to our knowledge, no work has been published systematically characterizing the responses of the Aag-2 cell line, against different kinds of pathogens and comparison of the response to the one exhibited by whole mosquitoes, in this way “validating” this cell line as a tool for immune investigations. For this reason, in this study we characterized the gene expression profile of the Aag-2 *Ae. aegypti* cell line in response to different kinds of immune challenges, such as Gram negative and positive bacteria, fungi and viruses. Our results show that the Aag-2 lineage is immunocompetent and also that the immune responses elicited closely resemble the responses described for whole *Ae. aegypti* mosquitoes.

Methods

Cell Culture

Aedes aegypti Aag-2 cells were maintained at 28 °C in Schneider's *Drosophila* medium with L-glutamine (Gibco) supplemented with 10% Fetal Bovine Serum (FBS) (Cultilab) and were passaged at a 1:10 dilution every 4–5 days. For this passage, cells were released from the culture flask with 0.25% trypsin solution (Gibco). Once in solution, cells were transferred to well plates for the indicated treatments or to another culture flask for maintenance.

Treatments and Viral Infection

Cells were seeded in a 12-well plate (TPP- Tecno Plastic Products) at 80-90% of confluence, approximately 2×10^5 cells per well and incubated overnight for adherence to occur. For bacterial challenge, cells were incubated with two different heat killed bacteria as previously described [34]: *Micrococcus luteus*, a Gram positive bacteria and *Enterobacter cloacae*, a Gram negative bacteria. Aag-2 cells were incubated with 100 bacteria per cell (10^7

bacteria/ well) or 1000 bacteria per cell (10^8 bacteria/ well). Cells were also stimulated with Zymosan A (Sigma-Aldrich), a yeast cell wall sugar that consists of protein-carbohydrate complexes. Two different concentrations of Zymosan A (0.5 mg/mL and 1 mg/mL) were used in the assays as previously described [36]. Before use, Zymosan A suspension was autoclaved to, eventually, eliminate any contaminant live yeast. The incubation with both bacteria and Zymosan A lasted 24 hours. For viral infection, cells were infected with Sindbis virus using a MOI (Multiplicity of Infection) of 10, as previously described [37]. Cells were incubated in the presence or absence of virus in Schneider's *Drosophila* medium with L-glutamine (Gibco) without FBS for 1 hour at 37°C. After that, cells were washed twice with phosphate-buffered saline (PBS) and medium supplemented with 10% SFB was replaced in all wells. The incubation lasted 24 hours and the cells were maintained at 28°C. Each figure represents at least five biological replicates.

RNA Extraction, cDNA synthesis and Quantitative PCR

Total RNA from cells in all conditions was extracted using the TRIZOL reagent (Invitrogen) following the manufacturer's instructions. RNA was treated with DNase I (Fermentas) and first-strand cDNA synthesis was carried out using High-Capacity cDNA Reverse transcription kit (Applied Biosystems). The efficiency of the experimental set for each gene was tested with serial dilutions of cDNA and was only used if the resultant efficiency was $\geq 90\%$. Each PCR reaction (15 μ L volume) contained diluted cDNA, 2 \times Power SYBR Green Master Mix (Applied Biosystems) and 300 nM of forward and reverse primers. Quantitative PCR was performed in a 7500 Real Time PCR System (Applied Biosystems) using Applied Biosystems recommended qPCR conditions (20 seconds at 95°C followed by 40 cycles of 95°C for 1 second and 20 seconds at 60°C followed by a melting curve to assure a simple product was amplified). The comparative $\Delta\Delta$ Ct method was used to evaluate changes in gene expression levels and all standard errors were calculated based on Δ Ct as described in Applied Biosystems User Bulletin #2 (http://www3.appliedbiosystems.com/cms/groups/mcb_support/documents/generaldocuments/cms_040980.pdf). The *Ae. aegypti* ribosomal protein 49 gene RP-49 was used as endogenous control (accession number AAT45939), based on previous data [38]. Primer sequences used are described on Table 1. Each figure represents at least five biological replicates with three technical replicates for each sample.

Phagocytosis assay

Cells were seeded in a 24-well plate with a glass slide at 60% of confluence. The phagocytosis assay was

performed as in Mizutani *et al.* (2003) [19]. The medium was removed and cells were incubated with 4×10^7 /mL 1.0 μ m fluorescent red latex beads carboxylate-modified polystyrene (Sigma Aldrich) per well in Mg^{2+} -free Hank's solution (Sigma Aldrich) and incubated for 1 hour at 28°C or 4°C (control). After this period, cells were washed with phosphate-buffered saline (PBS) and then fixed with 3.7% formaldehyde for 30 minutes. After fixation, 0.4% Trypan blue solution (Sigma Aldrich) in phosphate-buffered saline (PBS) was added to quench the fluorescence from the extracellular beads. Cells were observed in a fluorescence microscope Zeiss Axioskop 40 with an Axiocam MRC5 using a Zeiss-15 filter set (excitation BP 546–612; beam splitter FT 580; emission LP 590). Differential interference contrast (DIC) images were acquired with a Zeiss AxioObserver, which was also used for some fluorescence images, with Zeiss-15 filter set (excitation BP 546–612; beam splitter FT 580; emission LP 590) for fluorescent red latex beads. Comparison of fluorescence levels among distinct images was performed under identical conditions, using the same objectives, microscopes and similar exposure times in each experiment.

Statistical analysis

Quantitative PCRs were statistically analyzed by ANOVA followed by Dunnett's multiple comparison test. Analyses were performed on Δ Ct data, before normalization, using GraphPad Prism statistical software package (Prism 5.01; GraphPad Software, Inc., San Diego, CA). Asterisks indicate significant differences (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

Results

Cells were stimulated with two concentrations of heat-killed Gram negative (*Enterobacter cloacae*) and Gram positive (*Micrococcus luteus*) bacteria and with Zymosan A, a glucan from yeast cell walls, or infected with Sindbis virus.

Toll pathway

When cells were exposed to either Gram positive or Gram negative bacteria or to Zymosan, the genes from the Toll pathway presented a bimodal expression profile. The transcription factor REL 1 (Figure 1A) and the adaptor protein Myd88 (Figure 1B) did not change expression in comparison to control cells after any of the treatments, although a trend towards increase can be observed for REL 1 upon incubation of the cells with all immune elicitors.

The homolog of the NF- κ B inhibitor (I κ B), cactus, on the other hand, was significantly induced when cells were incubated with Zymosan (Figure 1C). The same profile was observed when cells were stimulated with

Table 1 Primers used in this study

Accession number	Gene	Primer sequence	Melting temperature (°C)	R ² efficiency Curve	Primer efficiency
AAEL007696	aaREL 1 For	GACTCGTCGGAGCTGAAATC	81.1	0.9989	1.1516
	aaREL 1 Rev	CGGTTTGTTTCAGGTTGTTGA			
AAEL007624	aaREL 2 For	TCTGTGCGGCAGATGAAGTGA	79.7	0.9999	1.1144
	aaREL 2 Rev	GCACTGGAATGGAGAATCAAA			
AAEL000709	Cactus For	TCTTGCGTTGAAGTGAGTGG	79.2	0.993	1.034
	Cactus Rev	GACCCCTCTGAAAGGGAAAGG			
AAEL003841	Defensin For	GATTTCGGCGTTGGTGATAGT	81.9	0.9992	0.9955
	Defensin Rev	TTATTCAATTCGGGCAGACG			
AAEL010083	IMD For	TCGTCAAACCTCGGTTTTCCT	78.9	0.9925	1.1068
	IMD Rev	TGGCGGAGTTGAAGGTAAAG			
AAEL007768	Myd88 For	CGATGCGTTTCATTTTGTGTTG	76.8	0.9889	1.03
	Myd88 Rev	CACCGCTCAGAAATCAGCTT			
AAT45939	RP49 For	GCTATGACAAGCTTGCCCCCA	83.7	0.9879	1.2002
	RP49 Rev	TCATCAGCACCTCCAGCT			
AAEL007765	Serpin For	ACGTGATGGATTGGATGGAG	79.2	0.9995	1.0149
	Serpin Rev	GTGCCTGCACTTGTTTCTGA			
AAEL009692	STAT For	CACACAAAAGGACGAAGCA	75.7	1	1.1797
	STAT Rev	TCCAGTTCCCTTAAAGCTCA			
AAEL001794	TEP For	ATTTTGTGACGGCTTTTGTGG	78.9	0.9992	1.0604
	TEP Rev	TGGATTACTTGCCCCACTTC			

the Gram positive bacteria or with high amounts of Gram negative bacteria (Figure 1C). An effector molecule of the Toll pathway, a serine protease inhibitor, serpin 27A, was also overexpressed in response to the two types of bacteria in both concentrations (Figure 1D). However, in cells incubated with Zymosan, the levels of serpin 27A mRNA were elevated only upon incubation of the cell with the smallest amount (Figure 1D).

When cells were infected with the alphavirus Sindbis, any significant increase in the expression profile of the analyzed Toll pathway related genes was observed. In fact, cactus and serpin presented a decrease in mRNA levels upon infection of the cells (Figure 2A and 2B).

IMD Pathway

REL 2, the NF-κB homologue transcription factor of IMD pathway, showed a 3 to 5 fold expression increase in response to incubations of Aag-2 cells with all concentrations of Zymosan, Gram negative and Gram positive bacteria (Figure 3A). The adaptor protein IMD, showed the same pattern observed for Myd88, the adaptor protein of the Toll pathway, having its expression unaltered in response to all stimuli (Figure 3B). One of the IMD pathway effectors, defensin was highly up-regulated in response to both Gram positive or negative bacteria and Zymosan (Figure 3C). Interestingly, the

level of defensin up-regulation was higher than any other molecule analyzed.

The IMD pathway showed any change in gene expression in cells infected with Sindbis virus (Figure 2C).

Jak/STAT Pathway

In the Jak/STAT pathway, transcription factor STAT did not show significant changes in response to Gram positive or negative bacteria or Zymosan at any concentration tested (Figure 4A). A thiol-ester motif-containing protein, TEP [39] was up-regulated in response to Gram negative and Gram positive bacteria (Figure 4B).

When the cells were infected with Sindbis virus, the transcription factor STAT was the only molecule analyzed that was significantly up-regulated (Figure 2D). However, the expression of TEP showed no changes in infected cells (Figure 2D).

Phagocytosis

An important characteristic of immune competent cells is the capacity to phagocytose foreign bodies such as bacteria or fungi. To evaluate if the cell line Aag-2, besides presenting a similar expression profile to immune cells, like hemocytes, was capable of phagocytosing microorganisms, cells were maintained in culture, incubated with fluorescent latex beads for 1 hour and observed in an epifluorescence microscope. Latex beads




Figure 1 Expression of Toll pathway marker genes after exposure of *Aedes aegypti* Aag-2 cells to Gram-positive (*M. luteus*) or negative (*E. cloacae*) bacteria or to Zymosan. Cells were incubated with the indicated amounts of the indicated stimuli for 24 hours and the expression of REL 1 (A), MyD88 (B), cactus (C) or serpin (D) was analyzed by qPCR using RP-49 gene as endogenous control. Each bar represents the relative expression and standard error of the analyzed genes, calculated as described in Methods section.

could be observed in the cytoplasm of virtually all cells in the well (Figure 5A-C), confirming the ability of this cell to phagocytose. As control, cells were incubated under the same conditions at 4°C to inhibit phagocytosis (Figure 5D-F). Incubations with Trypan Blue were made after the fixation step to quench the fluorescence from extracellular beads.

Discussion

Recently, insect cell lines have proven to be highly useful for such studies, since they are easy to handle and grow and typically produce homogeneous and reliable results.

In 1968, Peleg developed an *Ae. aegypti* embryonic cell line [31], Aag-2. Curiously for almost 30 years this lineage was neglected until Gao *et al.* (1999) [34] published a paper describing the secretion of a defensin by this cell in response to heat-killed Gram positive bacteria. After that, several groups started using Aag-2 cells and it has now become clear that this lineage has several advantages over C6/36 as a model cell for the genus *Aedes*. One important point is that, unlike *Ae. albopictus*, *Ae. aegypti* has a sequenced genome [14]. This is an extremely important characteristic of this cell since

having a genome makes it tremendously easier to identify genes of interest and use conventional methodologies to study gene expression, such as qPCR, but also allows the use of high-throughput research methodologies, such as microarrays [32]. Unlike C6/36, Aag-2 has a functional RNAi pathway [40]. This fact not only makes this cell a more accurate model to study viral infection but also allows the use of reversal genetic strategies, especially RNAi based approaches in this cell.

When the cells were exposed to Gram positive or negative bacteria or Zymosan, expression of cactus and serpin 27A were significantly increased. These results are consistent with an activation of the Toll pathway, since the expression of cactus is increased upon Toll pathway activation [41], and in whole mosquitoes, the expression of serpin 27A is responsive to the Toll pathway and is totally abolished upon knock down of REL 1 [42]. Interestingly, we also observed an activation of this pathway when the cells were incubated with Gram negative bacteria (*E. cloacae*), as seen in whole mosquitoes challenged with the same bacterium [35]. This finding reinforces the idea that Aag-2 immune responses are very similar to the mosquito.




Figure 2 Expression of Toll, IMD and Jak/STAT pathway marker genes after exposure of *Aedes aegypti* Aag-2 cells Sindbis virus. Cells were infected with Sindbis virus using a MOI of 10. After 24 hour of incubation at 28°C, the expression marker genes from the Toll (A,B), IMD(C) or Jak/STAT (D) pathways was analyzed by qPCR using RP49 gene as endogenous control. Each bar represents the relative expression and standard error of the analyzed genes, calculated as described in Methods section.

With the exception of the adapter protein IMD, all other IMD related genes investigated were induced by the three stimuli used. In *D. melanogaster*, defensin expression is controlled mainly by the Toll pathway, but in *Ae. aegypti*, this gene is controlled by the IMD pathway [43,44]. Defensin was induced more than 100 fold upon incubation with *E. cloacae*.

These results are significantly different for *Drosophila*, where the IMD pathway is activated only for Gram negative bacteria and not Gram positive or fungi [45].

Nevertheless, the immune pattern exhibited by Aag-2 cell accords to the pattern observed for whole mosquitoes. In mosquitoes, Gram-positive bacteria are able to activate the IMD pathway and the peptides produced by this pathway, defensin among them, are able to efficiently impair the growth of Gram-positive bacteria and increase mosquito survival upon infection [44,46].

From the “classical” mosquito immune pathways, Jak/STAT is the pathway about which less is known. The *Aedes* Jak/STAT pathway can be activated by fungal and




Figure 3 Expression of IMD pathway marker genes after exposure of *Aedes aegypti* Aag-2 cells to Gram-positive (*M. luteus*) or negative (*E. cloacae*) bacteria or to Zymosan. Cells were incubated with the indicated amounts of the indicated stimuli for 24 hours and the expression of REL 2 (A), IMD (B) or defensin (C) was analyzed by qPCR using RP49 gene as endogenous control. Each bar represents the relative expression and standard error of the analyzed genes, calculated as described in Methods section.




Figure 4 Expression of Jak/STAT pathway marker genes after exposure of *Aedes aegypti* Aag-2 cells to Gram-positive (*M. luteus*) or negative (*E. cloacae*) bacteria or to Zymosan. Cells were incubated with the indicated amounts of the indicated stimuli for 24 hours and the expression of STAT (A) or TEP (B) was analyzed by qPCR using RP49 gene as endogenous control. Each bar represents the relative expression and standard error of the analyzed genes, calculated as described in Methods section.

viral infections [39,47]. Here, we analyze two genes from this pathway, the transcription factor STAT and TEP, a gene shown to be under control of the Jak/STAT pathway [39]. STAT expression was not altered when incubated with bacteria or Zymosan. This is not surprising since the increase of STAT protein levels is not necessarily required for the activation of the pathway. TEP expression significantly increased in response to all these challenges, suggesting that all of them are able to activate Jak/STAT pathway. Again, this is in accordance with the data available in the literature for mosquitoes. *Beauveria bassiana*, an entomopathogenic, fungus activates Jak/STAT pathway [47]. Bartholomay *et al.* (2007) [48] showed that hemocytes of mosquitoes infected with a Gram negative (*E. coli*) or Gram positive (*M. luteus*) bacteria present significant increases in TEP expression.

When Aag-2 cells were infected with Sindbis virus, from all the genes of the three pathways we assayed, only STAT was significantly up-regulated. Although the mechanisms employed by mosquitoes to fight against flavivirus, such as dengue, have recently been revealed, with Toll and Jak/STAT, but not IMD, playing important roles in the control of virus titers [10,39], little is known about how mosquitoes control alphavirus infections. Sanders *et al.* (2005) [49] in a microarray study of *Aedes* infected with Sindbis virus, identified a transient increase in REL 1 expression only in the first day after infection. Curiously, no other gene from the Toll pathway was up-regulated in the array. On the other hand, depending on the Sindbis or *Aedes* strains used in this assay, no gene from the Toll pathway could be up-regulated in Sanders *et al.* (2005) [49] assay (Gill S,




Figure 5 *Aedes aegypti* Aag-2 cells are able to phagocytose. Aag-2 cells were incubated with 2×10^7 1.0 μ m fluorescent red latex beads (Sigma Aldrich) for 1 hour at 28°C (A-C) or 4°C (D-F, control). After this period, cells were fixed with formaldehyde 3.7% for 30 minutes and 0.4% Trypan blue solution in phosphate-buffered saline (PBS) and observed under a fluorescence microscope. (A and D) Differential interference contrast (DIC) images. (B and E) Fluorescence image. (C and F) Merge. Magnification 100X.

personal communication). In fact, we show that when Aag-2 cells were infected with Sindbis virus, no genes from the Toll pathway were up-regulated. On the contrary, one gene from the Toll pathway, cactus, was actually down-regulated.

The observed up-regulation of STAT upon Sindbis infection, could be related with the known involvement of this pathway in the control of viral infections [4,10]. Unfortunately, the lack of papers describing immune aspects, other than RNAi, of *Aedes*-Sindbis interaction makes it difficult to understand if the increased expression of STAT exhibited by Sindbis infected Aag-2 is also a characteristic of infected mosquitoes. Also noteworthy is the fact that, although STAT is up-regulated, there is no increase in TEP expression. This probably points to the fact that, upon Sindbis infection, only a subset of Jak/STAT regulated genes is activated.

Recent studies have unequivocally shown that a phagocytic response is as important as the humoral one for insects to fight pathogens [50]. Besides, the entrance of many viruses such as Dengue into host cell is dependent on active phagocytosis [51]. Our results show that, like mosquito hemocytes and other insect cultured cells, when exposed to latex beads Aag-2 efficiently phagocytes these bodies, revealing one more characteristic shared with other insect immunocompetent cells.

Conclusions

After examining Aag-2 cells immune responses against Gram-positive and negative bacteria, fungi and Sindbis virus, besides its capacity to phagocyte strange bodies, it is our conclusion that this cell lineage responds to these stimuli in a very similar way to that described for whole mosquitoes and constitutes a good model for insect immune studies.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Conceived and designed the experiments: ABBF MCLNS MHFS. Performed the experiments: ABBF. Analyzed the data: ABBF MCLNS MHFS. Contributed reagents/materials/analysis tools: MHFS. Wrote the paper: ABBF MHFS. All reviewers read and approved the final version of the manuscript.

Acknowledgements

We thank all of the members of the Laboratory of Biochemistry of Hematophagous Arthropods, especially Katia Anastácio Laia for rearing insects, S.R. Cássia for technical assistance. Dr. Clarissa Maia Monteiro for the fluorescent latex beads and Dr. Felipe Almeida Dias for *E. cloacae* and *M. luteus* used in experiments.

Author details

¹Laboratório de Bioquímica de Artrópodes Hematófagos, Instituto de Bioquímica Médica, Programa de Biologia Molecular e Biotecnologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil. ²Instituto Nacional de Ciência e Tecnologia em Entomologia Molecular (INCT-EM), Rio de Janeiro, Brazil. ³Trifermed, ESADECREAPOLIS innovation lab, Barcelona, Spain.

Received: 13 January 2012 Accepted: 24 July 2012

Published: 24 July 2012

References

1. *Dengue and haemorrhagic fever*; <http://www.who.int/mediacentre/factsheets/fs117/en/>.
2. WHO: Test procedures for insecticide resistance monitoring in malaria vectors, bio-efficacy and persistence of insecticides on treated surfaces. In *Book Test procedures for insecticide resistance monitoring in malaria vectors, bio-efficacy and persistence of insecticides on treated surfaces*. City: World Health Organization; 1998:46.
3. De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B: The Toll and Imd pathways are the major regulators of the immune response in *Drosophila*. *EMBO J* 2002, **21**:2568–2579.
4. Dostert C, Jouanguy E, Irving P, Troxler L, Galiana-Arnoux D, Hetru C, Hoffmann JA, Imler JL: The Jak-STAT signaling pathway is required but not sufficient for the antiviral response of *Drosophila*. *Nat Immunol* 2005, **6**:946–953.
5. Tanji T: Activation of the *Drosophila* innate immunity by the toll and IMD pathways. *Seikagaku* 2006, **78**:413–417.
6. Tanji T, Hu X, Weber AN, Ip YT: Toll and IMD pathways synergistically activate an innate immune response in *Drosophila melanogaster*. *Mol Cell Biol* 2007, **27**:4578–4588.
7. Tanji T, Ip YT: Regulators of the Toll and Imd pathways in the *Drosophila* innate immune response. *Trends Immunol* 2005, **26**:193–198.
8. Tzou P, Reichhart JM, Lemaitre B: Constitutive expression of a single antimicrobial peptide can restore wild-type resistance to infection in immunodeficient *Drosophila* mutants. *Proc Natl Acad Sci U S A* 2002, **99**:2152–2157.
9. Hoffmann JA, Reichhart JM: *Drosophila* innate immunity: an evolutionary perspective. *Nat Immunol* 2002, **3**:121–126.
10. Xi Z, Ramirez JL, Dimopoulos G: The *Aedes aegypti* toll pathway controls dengue virus infection. *PLoS Pathog* 2008, **4**:e1000098.
11. Lemaitre B, Reichhart JM, Hoffmann JA: *Drosophila* host defense: differential induction of antimicrobial peptide genes after infection by various classes of microorganisms. *Proc Natl Acad Sci U S A* 1997, **94**:14614–14619.
12. Kim T, Kim YJ: Overview of innate immunity in *Drosophila*. *J Biochem Mol Biol* 2005, **38**:121–127.
13. Ferrandon D, Imler JL, Hetru C, Hoffmann JA: The *Drosophila* systemic immune response: sensing and signalling during bacterial and fungal infections. *Nat Rev Immunol* 2007, **7**:862–874.
14. Nene V, Wortman JR, Lawson D, Haas B, Kodira C, Tu ZJ, Loftus B, Xi Z, Megy K, Grabherr M, et al: Genome sequence of *Aedes aegypti*, a major arbovirus vector. *Science* 2007, **316**:1718–1723.
15. Waterhouse RM, Kriventseva EV, Meister S, Xi Z, Alvarez KS, Bartholomay LC, Barillas-Mury C, Bian G, Blandin S, Christensen BM, et al: Evolutionary dynamics of immune-related genes and pathways in disease-vector mosquitoes. *Science* 2007, **316**:1738–1743.
16. Grace TD: Establishment of four strains of cells from insect tissues grown in vitro. *Nature* 1962, **195**:788–789.
17. Sanchez-Vargas I, Travanty EA, Keene KM, Franz AW, Beaty BJ, Blair CD, Olson KE: RNA interference, arthropod-borne viruses, and mosquitoes. *Virus Res* 2004, **102**:65–74.
18. Levashina EA, Moita LF, Blandin S, Vriend G, Lagueux M, Kafatos FC: Conserved role of a complement-like protein in phagocytosis revealed by dsRNA knockout in cultured cells of the mosquito, *Anopheles gambiae*. *Cell* 2001, **104**:709–718.
19. Mizutani T, Kobayashi M, Eshita Y, Shirato K, Kimura T, Ako Y, Miyoshi H, Takasaki T, Kurane I, Kariwa H, et al: Involvement of the JNK-like protein of the *Aedes albopictus* mosquito cell line, C6/36, in phagocytosis, endocytosis and infection of West Nile virus. *Insect Mol Biol* 2003, **12**:491–499.
20. Blandin SA, Levashina EA: Phagocytosis in mosquito immune responses. *Immunol Rev* 2007, **219**:8–16.
21. Cherry S: Genomic RNAi screening in *Drosophila* S2 cells: what have we learned about host-pathogen interactions? *Curr Opin Microbiol* 2008, **11**:262–270.
22. Pitaluga AN, Mason PW, Traub-Cseko YM: Non-specific antiviral response detected in RNA-treated cultured cells of the sandfly, *Lutzomyia longipalpis*. *Dev Comp Immunol* 2008, **32**:191–197.

23. Hughes GL, Ren X, Ramirez JL, Sakamoto JM, Bailey JA, Jedlicka AE, Rasgon JL: **Wolbachia infections in *Anopheles gambiae* cells: transcriptomic characterization of a novel host-symbiont interaction.** *PLoS Pathog* 2011, **7**:e1001296.
24. Moon AE, Walker AJ, Goodbourn S: **Regulation of transcription of the *Aedes albopictus* cecropin A1 gene: A role for p38 mitogen-activated protein kinase.** *Insect Biochem Mol Biol* 2011, **41**:628–636.
25. Adelman ZN, Sanchez-Vargas I, Travanty EA, Carlson JO, Beaty BJ, Blair CD, Olson KE: **RNA silencing of dengue virus type 2 replication in transformed C6/36 mosquito cells transcribing an inverted-repeat RNA derived from the virus genome.** *J Virol* 2002, **76**:12925–12933.
26. Igarashi A: **Isolation of a Singh's *Aedes albopictus* cell clone sensitive to Dengue and Chikungunya viruses.** *J Gen Virol* 1978, **40**:531–544.
27. Miller ML, Brown DT: **Morphogenesis of Sindbis virus in three subclones of *Aedes albopictus* (mosquito) cells.** *J Virol* 1992, **66**:4180–4190.
28. Singh KR: **Cell cultures derived from larvae of *Aedes albopictus* (Skuse) *Aedes aegypti* (L).** *Curr Sci* 1967, **36**:506–508.
29. Cheng LL, Bartholomay LC, Olson KE, Lowenberger C, Vizioli J, Higgs S, Beaty BJ, Christensen BM: **Characterization of an endogenous gene expressed in *Aedes aegypti* using an orally infectious recombinant Sindbis virus.** *J Insect Sci* 2001, **1**:10.
30. Brackney DE, Scott JC, Sagawa F, Woodward JE, Miller NA, Schilkey FD, Mudge J, Wilusz J, Olson KE, Blair CD, Ebel GD: **C6/36 *Aedes albopictus* cells have a dysfunctional antiviral RNA interference response.** *PLoS Negl Trop Dis* 2010, **4**:e856.
31. Peleg J: **Growth of arboviruses in primary tissue culture of *Aedes aegypti* embryos.** *Am J Trop Med Hyg* 1968, **17**:219–223.
32. Sim S, Dimopoulos G: **Dengue virus inhibits immune responses in *Aedes aegypti* cells.** *PLoS One* 2010, **5**:e10678.
33. Fallon AM, Sun D: **Exploration of mosquito immunity using cells in culture.** *Insect Biochem Mol Biol* 2001, **31**:263–278.
34. Gao Y, Hernandez VP, Fallon AM: **Immunity proteins from mosquito cell lines include three defensin A isoforms from *Aedes aegypti* and a defensin D from *Aedes albopictus*.** *Insect Mol Biol* 1999, **8**:311–318.
35. Shin SW, Kokoza V, Bian G, Cheon HM, Kim YJ, Raikhel AS: **REL1, a homologue of *Drosophila* dorsal, regulates toll antifungal immune pathway in the female mosquito *Aedes aegypti*.** *J Biol Chem* 2005, **280**:16499–16507.
36. Sung HH, Chang HJ, Her CH, Chang JC, Song YL: **Phenoloxidase activity of hemocytes derived from *Panagaea monodon* and *Macrobrachium rosenbergii*.** *J Invertebr Pathol* 1998, **71**:26–33.
37. Mukherjee S, Hanley KA: **RNA interference modulates replication of dengue virus in *Drosophila melanogaster* cells.** *BMC Microbiol* 2010, **10**:127.
38. Gentile C, Lima JB, Peixoto AA: **Isolation of a fragment homologous to the rp49 constitutive gene of *Drosophila* in the Neotropical malaria vector *Anopheles aquasalis* (Diptera: Culicidae).** *Mem Inst Oswaldo Cruz* 2005, **100**:545–547.
39. Souza-Neto JA, Sim S, Dimopoulos G: **An evolutionary conserved function of the JAK-STAT pathway in anti-dengue defense.** *Proc Natl Acad Sci U S A* 2009, **106**:17841–17846.
40. Scott JC, Brackney DE, Campbell CL, Bondu-Hawkins V, Hjelle B, Ebel GD, Olson KE, Blair CD: **Comparison of dengue virus type 2-specific small RNAs from RNA interference-competent and -incompetent mosquito cells.** *PLoS Negl Trop Dis* 2010, **4**:e848.
41. Nicolas E, Reichhart JM, Hoffmann JA, Lemaître B: **In vivo regulation of the *IkappaB* homologue cactus during the immune response of *Drosophila*.** *J Biol Chem* 1998, **273**:10463–10469.
42. Bian G, Shin SW, Cheon HM, Kokoza V, Raikhel AS: **Transgenic alteration of Toll immune pathway in the female mosquito *Aedes aegypti*.** *Proc Natl Acad Sci U S A* 2005, **102**:13568–13573.
43. Shin SW, Kokoza V, Ahmed A, Raikhel AS: **Characterization of three alternatively spliced isoforms of the Rel/NF-kappa B transcription factor Relish from the mosquito *Aedes aegypti*.** *Proc Natl Acad Sci U S A* 2002, **99**:9978–9983.
44. Antonova Y, Alvarez KS, Kim YJ, Kokoza V, Raikhel AS: **The role of NF-kappaB factor REL2 in the *Aedes aegypti* immune response.** *Insect Biochem Mol Biol* 2009, **39**:303–314.
45. Lemaître B, Hoffmann J: **The host defense of *Drosophila melanogaster*.** *Annu Rev Immunol* 2007, **25**:697–743.
46. Meister S, Kanzok SM, Zheng XL, Luna C, Li TR, Hoa NT, Clayton JR, White KP, Kafatos FC, Christophides GK, Zheng L: **Immune signaling pathways regulating bacterial and malaria parasite infection of the mosquito *Anopheles gambiae*.** *Proc Natl Acad Sci U S A* 2005, **102**:11420–11425.
47. Dong Y, Morton JC Jr, Ramirez JL, Souza-Neto JA, Dimopoulos G: **The entomopathogenic fungus *Beauveria bassiana* activate toll and JAK-STAT pathway-controlled effector genes and anti-dengue activity in *Aedes aegypti*.** *Insect Biochem Mol Biol* 2012, **42**:126–132.
48. Bartholomay LC, Mayhew GF, Fuchs JF, Rocheleau TA, Erickson SM, Aliota MT, Christensen BM: **Profiling infection responses in the haemocytes of the mosquito, *Aedes aegypti*.** *Insect Mol Biol* 2007, **16**:761–776.
49. Sanders HR, Foy BD, Evans AM, Ross LS, Beaty BJ, Olson KE, Gill SS: **Sindbis virus induces transport processes and alters expression of innate immunity pathway genes in the midgut of the disease vector, *Aedes aegypti*.** *Insect Biochem Mol Biol* 2005, **35**:1293–1307.
50. Nehme NT, Quintin J, Cho JH, Lee J, Lafarge MC, Kocks C, Ferrandon D: **Relative roles of the cellular and humoral responses in the *Drosophila* host defense against three gram-positive bacterial infections.** *PLoS One* 2011, **6**:e14743.
51. Mercer J, Schelhaas M, Helenius A: **Virus entry by endocytosis.** *Annu Rev Biochem* 2010, **79**:803–833.

doi:10.1186/1756-3305-5-148

Cite this article as: Barletta et al.: Validation of *Aedes aegypti* Aag-2 cells as a model for insect immune studies. *Parasites & Vectors* 2012 **5**:148.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

